

PUBLIC

ANNUAL REPORT 2025

OUT-OF-HOSPITAL CARDIAC ARREST 2025

by Deutsches Reanimationsregister – German Resuscitation Registry (GRR)

Deutsches
Reanimationsregister



Note:

The German Resuscitation Registry (GRR) is an online database with a constantly growing number of data sets. Due to different evaluation dates, the figures given may differ from those in previously published reports.

The tenth public annual report on out-of-hospital cardiac arrest (OHCA) by the German Resuscitation Registry (GRR) contains the current data, facts and figures on resuscitation provided by the participants in the German Resuscitation Registry in 2025.

This public annual report on out-of-hospital cardiac arrest is addressed to all participants and the public in order to further develop resuscitation care in Germany, fully in line with the Bad Boll Resuscitation and Emergency Dialogues and the overall social responsibility [1–3].

Data records from 208 emergency medical services (EMS), representing a population of approx. 37 million inhabitants, are used as a foundation. Thus, reliable

information on the resuscitation incidence in Germany, but also on the care of these patients and the success of treatment can be obtained.

As a limitation, it is to be noted that due to the voluntary participation in the German Resuscitation Registry (GRR) the data analysed represent a sample and may not be fully representative. This brief report is only a partial summary of the recorded data. The presentation is based on the Utstein Report [4,5], the international standardized reporting format for out-of-hospital cardiac arrest. If simplified of 'CPR' is spoken, it is out-of-hospital (EMS) resuscitation procedures for OHCA of various suspected or confirmed causes.

In the following, the overall data from the German Resuscitation Registry (GRR) for the period 01.01.2025 up to and including 31.12.2025 were analyzed. In addition, the data of a reference group of 56 German emergency medical services from the same period were analyzed, which met the following inclusion criteria:

- Incidence of resuscitation >30/100,000 inhabitants and year
- ROSC (Return Of Spontaneous Circulation) <80%
- RACA (ROSC after Cardiac Arrest) score calculable >60%
- Percentage of documented follow-ups of at least 30%

By determining the reference data, the results become more accurate, and statements regarding the discharge rate and the neurological outcomes at discharge are also possible.

1. Fischer M, Wnent J, Gross B, Seewald S, Maurer H, Ramshorn-Zimmer AB et al: Qualitätsmanagement in der gesamten Reanimationsversorgung ist unerlässlich. These 9 der Bad Boller Reanimations- und Notfallgespräche 2023. *Anästh Intensivmed* 2023;64:523–527
2. Gräsner JT, Wnent J, Zwißler B, Beck G, Fischer M: 10. Bad Boller Reanimations- und Notfallgespräche – Zeit für ein Update. *Anästh Intensivmed* 2023;64:473–475
3. Wnent J, Geldner G, Werner C, Bottiger BW, Fischer M, Scholz J, et al: Bad Boller resuscitation talks: 10 basic ideas for 10,000 lives. *Anesthesiol Intensivmed Notfallmed Schmerzther* 2014;49(3):208
4. Perkins GD, Jacobs IG, Nadkarni VM, Berg RA, Bhanji F, Biarent D, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein Resuscitation Registry Templates for Out-of-Hospital Cardiac Arrest: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation* 2015;132:1286–1300. Epub 11. November 2014
5. Bray JE, Gräsner JT, Nolan JP, Iwami T, Ong MEH, Finn J, et al: Cardiac Arrest and Cardiopulmonary Resuscitation Outcome Reports: 2024 Update of the Utstein Out-of-Hospital Cardiac Arrest Registry Template. *Circulation* 2024;150:e203–e223



CONTENTS

Number of patients and emergency medical services	6
Incidence of out-of-hospital cardiac arrest	6
Age of patients	7
Gender distribution	7
CPR before EMS arrival	9
Sites where cardiac arrest occurs	10
Witnessed cardiac arrests	10
Telephone guided CPR	11
Time between alarm and arrival of the 1st vehicle	11
Suspected cause of cardiac arrest	12
First recorded ECG rhythm	13
Resuscitation procedures	14
Use of mechanical chest compression devices	14
Outcome of EMS treatment	15
In-hospital treatment	16
Summary	17
Authors	18

NUMBER OF PATIENTS AND EMERGENCY MEDICAL SERVICES

26,438

from 208 emergency medical services

In 2025, the total number of out-of-hospital cardiac arrests documented in the German Resuscitation Registry was 26,438 from 208 emergency medical services.

12,116

from 56 emergency medical services

The chosen reference group for this report meets the above mentioned inclusion criteria and comprises 12,116 patients from 56 EMS.

The reference group is highlighted in the report. This group is supplemented by the overall data. The overall data is indicated as such in this report.



INCIDENCE OF OUT-OF-HOSPITAL CARDIAC ARREST

In 2025, the incidence of CPR was 71.6 CPRs per 100,000 inhabitants per year in the overall data. Extrapolated to Germany's current population of 83.5 million, in 2025 approximately 59,800 patients were resuscitated by the emergency medical services after a sudden cardiac arrest.

At the reference sites, the incidence of death declarations, resuscitations, and performed resuscitation treatments was significantly higher than in the overall group. Therefore, underreporting in the overall data cannot be ruled out.

declaration of death and resuscitation

154.6 | 137.6
per 100,000 inhabitants per year | overall data

CPR by EMS

77.5 | 71.6
per 100,000 inhabitants per year | overall data

AGE OF PATIENTS

The average patient is an approx. 69-year-old male. Very young patients - younger than 18 years - were rarely resuscitated. As in the previous year, just over 45% of those affected suffered an out-of-hospital cardiac arrest with resuscitation while of working age (under 70). The number of patients aged over 80 years has increased over time; their proportion is now 32.1% (2014: 27.7%).



GENDER DISTRIBUTION







CPR BEFORE EMS ARRIVAL

First Responder

In Germany, first responders are voluntary units that are not part of the regular EMS but are used by the dispatch centres to bridge the interval until the EMS arrives. This includes people who work in BOS (authorities and organisations with security tasks) (e.g. fire brigade). In the future, the data of first responders and activated bystanders will be recorded separately in accordance with the current Utstein report.

Activated Bystander

Activated bystanders are individuals who are alerted, for example, via apps and who bridge the interval until the EMS arrives. These can be trained first responders as well as medical professionals who are off duty and, by definition, not on call.

Bystander

Non-alert persons who happen to be at the scene of an emergency by chance and bridge the time between the incident and the arrival of the EMS, thus shortening the non-ambulance interval.

For years, public campaigns have aimed to draw attention to sudden cardiac death and possible CPR by bystanders. Furthermore, the guidelines on CPR recommend telephone guided CPR by dispatch centers.

The 2025 annual report shows a bystander CPR rate of 53.5% in the reference group. In 6.8% of the cases, first responders carried out CPR before the EMS arrived, in 2.8% of the cases activated bystanders started CPR.

Telephone guided CPR was 38.4% in the reference group.

total

63.1%

60.9%
overall data

by bystanders

53.5% | 51.9%
overall data


by first responders

6.8% | 7.0%
overall data

by activated bystanders


2.8% | 2.0%
overall data

SITES WHERE CARDIAC ARREST OCCURS




at home

68.7%
overall data



in public

15.9%
overall data



other sites

15.4%
overall data

WITNESSED CARDIAC ARRESTS

overall observed cardiac arrest



56.3%
overall data

by bystanders

42.4%

41.3%
overall data

by first responders


1.1%

1.5%
overall data

by EMS

14.0%

13.5%
overall data



TELEPHONE GUIDED CPR

The vast majority of resuscitations in 2025 also took place in private homes.

38.4%

37.3%
overall data

In recent years, we have seen a steady increase in the number of cases supported by telephone guided CPR. However, there is still considerable potential for growth [6].



TIME BETWEEN ALARM AND ARRIVAL OF THE 1ST VEHICLE

The average time between receiving the alarm and arriving at the scene with the first EMS unit (RTW or NEF) was **6 minutes and 40 seconds**.

06:40 ± 03:43


07:07 ± 04:14
overall data



RTW = Rettungswagen, ALS-ambulance **NEF** = Notarzteinsatzfahrzeug, Emergency physician's car

SUSPECTED CAUSE OF CARDIAC ARREST

Information on causes of cardiac arrest is based on the suspected diagnoses of the emergency physicians. Of course, this information may differ from the exact diagnoses.



FIRST RECORDED ECG RHYTHM

shockable (VF/VT)

22.9%

21.1%
overall data

VF = ventricular fibrillation
VT = ventricular tachycardia

non-shockable

77.2%

78.9%
overall data

asystole

55.3% | **54.8%**
overall data

pulseless electrical activity

23.6% | **22.4%**
overall data

cardiac cause


52.1% | 52.2%
overall data



RESUSCITATION PROCEDURES

Compared to 2024, significant changes can be seen in the area of airway management. Endotracheal intubation with and without a video laryngoscope decreased. This is a problematic development, as it has been shown in Germany and Austria [7] that endotracheal intubation increases the discharge rate compared to using supraglottic airway alone.

The other resuscitation measures remained largely unchanged from the previous year. In particular, the use of intraosseous access remains too high at 23.3%, although the new ERC-guidelines published in 2025 clearly indicate that intravenous access should be favored in resuscitation [8,9].



USE OF MECHANICAL CHEST COMPRESSION DEVICES

13.2% | 12.4% overall data

OUTCOME OF TREATMENT

In 2025 31.9 % (reference group) respectively 30.1 % (overall data) of patients reached the hospital with spontaneous circulation.

Standardized to 100,000 inhabitants per year, around nine patients per 100,000 inhabitants were discharged from hospital alive.

	ROSC	41.3%	overall data 39.7%
ROSC on admission to hospital		31.9%	30.1%
24-hour survival		20.1%	
discharged alive from hospital		10.9%	

7. Sulzgruber P, et al: The impact of airway strategy on the patient outcome after out-of-hospital cardiac arrest: A propensity score matched analysis. Eur Heart J Acute Cardiovasc Care, 2017;2048872617731894

8. Soar J, et al: European Resuscitation Council Guidelines 2021: Adult advanced life support. Resuscitation 2021;161:115–151


9. Soar J, et al: European Resuscitation Council Guidelines 2025: Adult advanced life support. Resuscitation 2025;215

IN-HOSPITAL TREATMENT

Hospital treatment has a significant impact on survival after successful resuscitation. Current guidelines recommend coronary angiography and intervention in particular, as well as temperature management [10]. Coronary angiography rates declined significantly in 2025 compared to 2024, while continuous temperature management remained stable at approximately 20 percent. Since 2021, the ERC guidelines have recommended temperature management for every initial rhythm and regardless of whether the cardiac arrest occurred inside or outside the hospital [11]. It remains to be seen whether the rate of interventions will increase and whether the guidelines will be implemented accordingly.

10. Nolan JP, et al: European Resuscitation Council and European Society of Intensive Care Medicine Guidelines 2025: Post-Resuscitation Care. Resuscitation 2025;215

11. Nolan JP, et al: European Resuscitation Council and European Society of Intensive Care Medicine guidelines 2021: post-resuscitation care. Intensive Care Med 2021;47(4):369–421



SUMMARY

The present annual report 'Out-of-hospital cardiac arrest 2025' by the German Resuscitation Registry (GRR) confirms the overall good quality of CPR care in the participating emergency medical services.

In 2025, the data from the resuscitation registry also documented deviations from guidelines and recommendations. For instance, in terms of response times, the target of 80% for reaching patients within 8 minutes of the emergency call was missed. According to current guidelines, there is still potential for improvement in areas such as telephone CPR, airway management, vascular access, the use of mechanical chest compression devices, and temperature management. A positive development worth noting is that more than half of those affected receive resuscitation before the EMS arrives.

We want to thank all participating emergency medical services. As of the cut-off date in 2025, data from 26,438 patients were recorded in the registry.

The German Resuscitation Registry was developed as a quality assurance tool for emergency medical services. In

addition to monthly and annual reports, the participants also receive access to online analyses in order to continuously benchmark, assess and improve the performance of their service - as required by the current resuscitation guidelines [12]. For example, the ROSC-after-Cardiac-Arrest (RACA) score [13] developed from the registry data is available in the online database for this purpose. The annual report of the German Resuscitation Registry on out-of-hospital cardiac arrest in 2025 provides information on the status and trends in resuscitation care in Germany and forms the basis for discussions.

Since cardiac arrest and resuscitation treatment are among the most time-critical and complex clinical conditions in emergency medicine, this result should be an incentive to further improve quality management and the overall quality of care in one's own emergency medical services.

The data on which this report is based and further information can be found in *Anästhesiologie und Intensivmedizin* (Anästh Intensivmed 2026;67:V1-V11. DOI: 10.19224/ai2026.1).

12. Greif R, et al: European Resuscitation Council Guidelines 2025: Executive Summary. *Resuscitation* 2025;215

13. Gräsner JT, et al: ROSC after cardiac arrest – the RACA score to predict outcome after out-of-hospital cardiac arrest. *Eur Heart J* 2011;32(13):1649–1656

AUTHORS



Prof. Dr. Matthias Fischer

- » University Medical Center Schleswig-Holstein, Institute for Emergency Medicine, Kiel
- » » agswm - Arbeitsgemeinschaft Südwestdeutscher Notärzte e.V., Filderstadt



PD Dr. Jan Wnent

- » University Medical Center Schleswig-Holstein, Institute for Emergency Medicine, Kiel
- » University Medical Center Schleswig-Holstein, Department of Anaesthesiology and Intensive Care Medicine, Campus Kiel
- » University of Namibia, School of Medicine, Windhoek, Namibia



Prof. Dr. Jan-Thorsten Gräsner

- » University Medical Center Schleswig-Holstein, Institute for Emergency Medicine, Kiel
- » University Medical Center Schleswig-Holstein, Department of Anaesthesiology and Intensive Care Medicine, Campus Kiel



PD Dr. Stephan Seewald

- » University Medical Center Schleswig-Holstein, Department of Anaesthesiology and Intensive Care Medicine, Campus Kiel



Lisa Rück

- » University Medical Center Schleswig-Holstein, Institute for Emergency Medicine, Kiel



Hanna Hoffmann

» University Medical Center Schleswig-Holstein, Institute for Emergency Medicine, Kiel



Prof. Dr. Berthold Bein

» Asklepios Klinik St. Georg, Department of Anaesthesiology and Intensive Care Medicine, Hamburg



Dr. Alexandra Ramshorn-Zimmer

» University of Leipzig Medical Center, Medical Management Division,
Department of Clinical Process Management



Prof. Dr. Andreas Bohn

» University Hospital Münster, Institute for Tele-Intensive and Preclinical Tele-Emergency
Medicine at the Department for Anaesthesiology, Surgical Intensive Care Medicine
and Pain Therapy

» City of Münster, Fire Department, Medical Director of the Emergency Medical Services

... and the participating Emergency Medical Services in the German Resuscitation Registry (GRR)

www.reanimationsregister.de

Deutsche Gesellschaft für Anästhesiologie
und Intensivmedizin e.V. (DGAI)

Neuwieder Straße 9
90411 Nürnberg

kontakt@reanimationsregister.de

Photo credits:
© by University Medical
Center Schleswig-Holstein

Deutsches Reanimationsregister –
German Resuscitation Registry (GRR)

Copyright © by Deutsches Reanimationsregister 2026

Suggested citation:

Fischer, M., Whent, J., Gräsner, J.-T., Seewald, S., Rück, L., Hoffmann, H.,
Bein, B., Ramshorn-Zimmer, A., Bohn, A. & the participating Emergency
Medical Services in the German Resuscitation Registry (GRR). (2026).
*Public Annual Report 2025 on Out-of-Hospital Cardiac Arrest 2025 by
Deutsches Reanimationsregister - German Resuscitation Registry (GRR).*
www.reanimationsregister.de/berichte.html